

# PRO microneedling

## contraindications

This list of contraindications are specific to PRO Microneedling.

When determining whether a treatment is suitable or not, it is important to use professional judgement and consider standard service or product contraindications.

### **ABSOLUTE**

- x Cancer
- x Malignant tumours
- x Pregnancy or breastfeeding
- x Any contagious or transmittable diseases
- x Chronic inflammatory skin conditions
- x Keloid or excessive hypertrophic scarring
- x Haemophilia or other cardiac abnormalities or bleeding disorders
- x Autoimmune disorders or immunosuppression
- x Dysfunction of the nervous system
- x Tuberculosis or liver function issues
- x Epilepsy
- x Disorders of collagen formation such as **scleroderma**
- x Medications such as those for blood disorders and steroids
- x Blood thinning medication
- x Prescription retinoids such as isotretinoin
- x Allergy to stainless steel

### **RELATIVE**

- x Health or medical concerns that require medical approval such as irregular blood pressure
- x Unwell within the last 2 weeks with a cold or flu virus
- x Medication such as isotretinoin within 6 months to 1 year from last dose
- x Medication that would affect the characteristics of the skin within 2 months
- x Aspirin
- x Facelift or eyelid surgery within the past year
- x Major operation scars less than 2 years old on the treated area
- x Dermabrasion, remodeling, deep chemical peels or surgical procedure with the last 3 months
- x Professional resurfacing, dermaplaning or superficial chemical peel within the last 14-28 days
- x IPL or laser procedures within the last 14 days
- x Excessively tanned or sunburnt skin from uv exposure or tanning creams within the last 2 weeks
- x Dermal filler administration within 14 days
- x Facial waxing, depilatory creams or electrolysis within the last 5 days
- x Alcohol or caffeine consumption within the last 24-48 hours
- x Facial shaving within the last 24 hours
- x Retinoid use within 3 days
- x Chronic stress

# PRO microneedling

## contraindications

### LOCAL

- x Facial melanosia
- x Active acne, actinic solar keratosis or rosacea
- x Herpes simplex
- x Raised moles, warts, skin lesions and scarring
- x Open cuts, wounds, abrasions and bruises.
- x Capillary ectasia
- x Botox, collagen, fat or other methods of augmentation within 4 weeks
- x Undiagnosed lumps, inflammation or swelling
- x Mild or occasional eczema, psoriasis or dermatitis
- x Piercings in the area

\*Always check manufacturer guidance for individual devices

Please note:

If medical status has changed or vaccination has taken place you are required to **update patch testing** before treating with Pro Power Peel

Avoid patch testing within one week of vaccination or experiencing Covid, as sensitivity may be heightened.

